arXiv:1007.0646v2 [math.AG] 27 Jul 2010

$\mathbb{G}_a^M$ DEGENERATION OF FLAG VARIETIES

EVGENY FEIGIN

Abstract. Let $\mathcal{F}_\lambda$ be a generalized flag variety of a simple Lie group G embedded into the projectivization of an irreducible G-module V_λ. We define a flat degeneration $\mathcal{F}_\lambda^a$, which is a $\mathbb{G}_a^M$ variety. Moreover, there exists a larger group G^a acting on $\mathcal{F}_\lambda^a$, which is a degeneration of the group G. The group G^a contains $\mathbb{G}_a^M$ as a normal subgroup. If G is of type A, then the degenerate flag varieties can be embedded into the product of Grassmanians and thus to the product of projective spaces. The defining ideal of $\mathcal{F}_\lambda^a$ is generated by the set of degenerate Plücker relations. We prove that the coordinate ring of $\mathcal{F}_\lambda^a$ is isomorphic to a direct sum of dual PBW-graded $\mathfrak{g}$-modules. We also prove that there exist bases in multi-homogeneous components of the coordinate rings, parametrized by the semistandard PBW-tableux, which are analogues of semistandard tableux.

Introduction

Let G be a simple Lie group with the Lie algebra $\mathfrak{g}$. The flag varieties associated with G are the quotients G/P by the parabolic subgroups (see e.g.[K1]). These varieties can be realized as G-orbits $G[v_\lambda] \hookrightarrow \mathbb{P}(V_\lambda)$ in the projectivization of irreducible G-modules V_λ with highest weight vector v_λ, $[v_\lambda] = \mathbb{C}v_\lambda$. We denote $\mathcal{F}_\lambda = G[v_\lambda]$.

In this paper we introduce a new family of varieties $\mathcal{F}_\lambda^a$, which are flat degenerations of $\mathcal{F}_\lambda$ (the superscript a is for abelian, see the explanations below). We note that in the literature there exists a degeneration of flag varieties into toric varieties (see [C], [GL], [L]). Our degenerate flags are $\mathbb{G}_a^M$ varieties, i.e. they are equipped with an action of the M-fold product $\mathbb{G}_a^M$ of the additive group of the field with an open orbit, where M is the dimension of a maximal unipotent subgroup of G.

The varieties $\mathcal{F}_\lambda^a$ are defined as follows. Let F_s, $s \geq 0$ be the PBW filtration on V_λ (see [Fe1], [Fe2], [FFoL], [K2])

$$F_s = \mathrm{span}\{x_1 \dots x_l v_\lambda : \ x_i \in \mathfrak{g}, l \leq s\}.$$

We define $V_\lambda^a = F_0 \oplus_{s\geq 0} F_{s+1}/F_s$. Let $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \mathfrak{n}^-$ be the Cartan decomposition. The space V_λ^a has a natural structure of a module over the degenerate algebra $\mathfrak{g}^a$. The algebra $\mathfrak{g}^a$ is isomorphic to $\mathfrak{g}$ as a vector space. It is a direct sum of two subalgebras: one is the Borel subalgebra $\mathfrak{b} = \mathfrak{n} \oplus \mathfrak{h}$ and the second is an abelian ideal $(\mathfrak{n}^-)^a$, which is an abelian subalgebra isomorphic to $\mathfrak{n}^-$ as a vector space. The corresponding Lie group G^a is a semidirect product $\mathbb{G}_a^M \rtimes B$ of the normal subgroup $\mathbb{G}_a^M$ ($M = \dim \mathfrak{n}$)

and the Borel subgroup B. We define the varieties $\mathcal{F}^a_\lambda$ as the closure of the $\mathbb{G}_a^M$-orbit of the highest weight vector:

$$\mathcal{F}_\lambda = \overline{\mathbb{G}^M[v_\lambda]} \hookrightarrow \mathbb{P}(V^a_\lambda).$$

We note that an affine space $\mathbb{G}_a^M[v_\lambda]$ does not coincide with the closure $\overline{\mathbb{G}_a^M[v_\lambda]}$, but form a dense open $\mathbb{G}_a^M$-orbit. Hence for all λ the varieties $\mathcal{F}_\lambda$ are the so-called $\mathbb{G}_a^M$ varieties [HT], [AS], [A].

We study the varieties $\mathcal{F}^a_\lambda$ in the case $\mathfrak{g} = \mathfrak{sl}_n$. In this case the varieties $\mathcal{F}_\lambda$ are isomorphic to partial flag varieties. In particular, the ones corresponding to fundamental weights are Grassmanians $Gr(d,n)$. There exist embeddings of partial flags into the product of projective spaces and the image is given by Plücker relations. These relations describe coordinate rings of flag varieties (see [Fu]). Since all fundamental weights are cominuscule in the $\mathfrak{sl}_n$ case, we have $\mathcal{F}^a_{\omega_d} \simeq \mathcal{F}_{\omega_d} \simeq Gr(d,n)$. The results of [FFoL] imply that each degenerate flag variety can be embedded into the product of Grassmanians and thus into the product of projective spaces. We show that this embedding can be described in terms of the explicit set of multi-homogeneous algebraic equations, which are obtained from the Plücker relations by certain degeneration. We prove that the degeneration $\mathcal{F} \to \mathcal{F}^a$ is flat. We also show that (as in the classical case) the coordinate ring of $\mathcal{F}^a_\lambda$ is isomorphic to a direct sum of certain dual modules $(V^a_\mu)^*$.

Our main technical tool is combinatorics of the semistandard PBW-tableaux. Recall that the usual semistandard tableaux serve as labeling set for bases of the coordinate rings of flag varieties. We construct another set (which we call the set of semistandard PBW-tableux) of Young diagrams filled by numbers such that these tableaux provide a basis in the coordinate rings of $\mathcal{F}^a_\lambda$ (and of $\mathcal{F}_\lambda$ as well).

Finally, we note that the case $\mathfrak{g} = \mathfrak{sl}_n$ is special, since all fundamental weights are cominuscule. This is important for our construction, since the operators from the nilpotent radical act as pairwise commuting operators in each module V_{ω_d} even before passing to $V^a_{\omega_d}$. This gives an identification $\mathcal{F}^a_{\omega_d} \simeq \mathcal{F}_{\omega_d} \simeq Gr(d,n)$ and simplifies the whole picture (see [A]). For other algebras there exist Grassmanians, which have to be degenerated. It is very interesting to study these degenerations as well as arbitrary degenerations $\mathcal{F}^a_\lambda$ for general $\mathfrak{g}$.

Our paper is organized as follows:
In Section 1 we recall notations and main facts about Lie algebras and flag varieties in type A.
In Section 2 we introduce the $\mathbb{G}_a^M$ degeneration of the generalized flag varieties for arbitrary simple Lie group G.
In Section 3 we state our main results and provide examples.
In Sections 4 and 5 we prove the statements from Section 3. Section 4 addresses the combinatorial questions and in Section 5 we prove algebro-geometric results.

1. The classical case

1.1. **Notations.** Let $\mathfrak{g}$ be a simple Lie algebra with the Cartan decomposition $\mathfrak{g}=\mathfrak{n}\oplus\mathfrak{h}\oplus\mathfrak{n}^-$ and the Borel subalgebra $\mathfrak{b}=\mathfrak{n}\oplus\mathfrak{h}$. Let ω_i, α_i, $i=1,\dots,l$ be fundamental weights and simple roots of $\mathfrak{g}$. Here $l=\dim\mathfrak{h}=\mathrm{rk}\ \mathfrak{g}$. Let Q and P be root and weight lattices in $\mathfrak{h}^*$: P is generated by ω_i and Q is generated by α_i. We set

$$Q_\pm=\bigoplus_{i=1}^l \pm\mathbb{Z}_{\ge0}\alpha_i,\ P_\pm=\bigoplus_{i=1}^l \pm\mathbb{Z}_{\ge0}\omega_i,\ \mathbb{Z}_{\ge0}=\{n\in\mathbb{Z}:\ n\ge0\}.$$

We denote by $(\cdot,\cdot)$ the Killing form on $\mathfrak{h}^*$. In particular, we have $(\alpha_i,\omega_j)=\delta_{i,j}$.

Consider the weight decomposition

$$\mathfrak{n}=\bigoplus_{\alpha\in Q_+}\mathfrak{n}_\alpha,\ \mathfrak{n}^-=\bigoplus_{\alpha\in Q_+}\mathfrak{n}^-_{-\alpha},$$

where $\mathfrak{n}_\alpha$ and $\mathfrak{n}^-_{-\alpha}$ are one-dimensional spaces spanned by elements e_α and f_α. One has

$$[h,e_\alpha]=\alpha(h)e_\alpha,\ [h,f_\alpha]=-\alpha(h)f_\alpha,\ h\in\mathfrak{h}.$$

We denote the elements e_{α_i} by e_i and f_{α_i} by f_i. Then e_α, f_α, $\alpha\in Q_+$ and $h_i=[e_i,f_i]$ form the Chevalley basis of $\mathfrak{g}$.

For any $\lambda=\sum_{i=1}^l m_i\omega_i\in P_+$ let V_λ be irreducible highest weight $\mathfrak{g}$-module with highest weight λ. Let $v_\lambda\in V_\lambda$ be a highest weight vector. Then one has

$$hv_\lambda=\lambda(h)v_\lambda\ \forall h\in\mathfrak{h},\quad \mathfrak{n}v_\lambda=0,\ \mathrm{U}(\mathfrak{n}^-)v_\lambda=V_\lambda.$$

Let G be a simple Lie group with the Lie algebra $\mathfrak{g}$. Let B, H, N and N^- be the Borel, Cartan and unipotent subgroups of G. The corresponding Lie algebras are $\mathfrak{b}$, $\mathfrak{h}$, $\mathfrak{n}$ and $\mathfrak{n}^-$. Each space V_λ, $\lambda\in P_+$ is equipped with the natural structure of G-module. Therefore G acts on the projectivization $\mathbb{P}(V_\lambda)$. The (generalized) flag variety $\mathcal{F}_\lambda\hookrightarrow\mathbb{P}(V_\lambda)$ is defined as the G-orbit of the line $\mathbb{C}v_\lambda$. Each variety $\mathcal{F}_\lambda$ is isomorphic to the quotient of G by the parabolic subgroup leaving the point $\mathbb{C}v_\lambda\in\mathbb{P}(V_\lambda)$ invariant.

1.2. **Type A case: Plücker relations, coordinate rings and semistandard tableaux.** In this subsection we recall the main ingredients of the theory of flag varieties in the case $\mathfrak{g}=\mathfrak{sl}_n$. Our main reference is [Fu] (see [K1] for more Lie-theoretic approach).

Let $1\le d_1<\dots<d_s\le n-1$ be a sequence of increasing numbers. Then for any positive integers $a_1,\dots,a_s$ the variety $\mathcal{F}_{a_1\omega_{d_1}+\dots+a_s\omega_{d_s}}$ is isomorphic to the partial flag variety

$$\mathcal{F}(d_1,\dots,d_s)=\{V_1\hookrightarrow V_2\hookrightarrow\dots\hookrightarrow V_s\hookrightarrow\mathbb{C}^n:\ \dim V_i=d_i\}.$$

In particular, if $s = 1$, then $\mathcal{F}(d)$ is the Grassmanian $Gr(d,n)$ and for $s = n-1$ $\mathcal{F}(1,\dots,n-1)$ is the variety of full flags. We recall that

$$V_{\omega_d} = \Lambda^d(V_{\omega_1}) = \Lambda^d(\mathbb{C}^n)$$

and the embedding $Gr(d,n) \hookrightarrow \mathbb{P}(\Lambda^d V_{\omega_1})$ is defined as follows: a subspace with a basis $w_1,\dots,w_d$ maps to $\mathbb{C}w_1 \wedge \dots \wedge w_d$. For general sequence $d_1,\dots,d_s$ one has embeddings:

$$\mathcal{F}(d_1,\dots,d_s) \hookrightarrow Gr(d_1,n) \times \dots \times Gr(d_s,n) \hookrightarrow \mathbb{P}(V_{\omega_{d_1}}) \times \dots \times \mathbb{P}(V_{\omega_{d_s}}).$$

The composition of these embeddings is called the Plücker embedding. The image is described explicitly in terms of Plücker relations. Namely, let $v_1,\dots,v_n$ be a basis of $\mathbb{C}^n = V_{\omega_1}$. Then one gets a basis v_J of V_{ω_d} $v_J = v_{j_1} \wedge \dots \wedge v_{j_d}$ labeled by sequences $J = (1 \le j_1 < j_2 < \dots < j_d \le n)$. Let $X_J \in V^*_{\omega_d}$ be the dual basis. We denote by the same symbols the coordinates of a vector $v \in V_{\omega_d}$: $X_J = X_J(v)$. The image of the embedding

$$\mathcal{F}(d_1,\dots,d_s) \hookrightarrow \times_{i=1}^s \mathbb{P}(V_{\omega_{d_i}})$$

is defined by the Plücker relations. These relations are labeled by a pair of numbers $p \ge q$, $p,q \in \{d_1,\dots,d_s\}$, by a number k, $1 \le k \le q$ and by a pair of sequences $L = (l_1,\dots,l_p)$, $J = (j_1,\dots,j_q)$, $1 \le l_\alpha, j_\beta \le n$. The corresponding relation is denoted by $R^k_{L,J}$ and is given by

$$R^k_{L,J} = X_L X_J - \sum_{1 \le r_1 < \dots < r_k \le p} X_{L'} X_{J'}, \tag{1.1}$$

where L',J' are obtained from L, J by interchanging k-tuples $(l_{r_1},\dots,l_{r_k})$ and $(j_1,\dots,j_k)$ in L and J respectively, i.e.

$$J' = (l_{r_1},\dots,l_{r_k}, j_{k+1},\dots,j_q),$$
$$L' = (l_1,\dots,l_{r_1-1}, j_1, l_{r_1+1},\dots,l_{r_2-1}, j_2,\dots,l_p).$$

We note that for any $\sigma \in S_d$ the equality

$$X_{j_{\sigma(1)},\dots,j_{\sigma(d)}} = (-1)^\sigma X_{j_1,\dots,j_d}$$

is assumed in (1.1). We denote the ideal generated by all $R^k_{L,J}$ by $I(d_1,\dots,d_s)$. An important fact is that this ideal is prime.

Recall that the coordinate ring of $\mathcal{F}(d_1,\dots,d_s)$ is the quotient ring

$$Q(d_1,\dots,d_s) = \mathbb{C}[X_{j_1,\dots,j_d}]/I(d_1,\dots,d_s),$$

where the variables $X_{j_1,\dots,j_d}$ are labeled by $d = d_1,\dots,d_s$ and $1 \le j_1 < \dots < j_d \le n$.

The ring $Q(d_1,\dots,d_s)$ has two other descriptions. First, recall that for any $\lambda,\mu \in P^+$ there exists an embedding of $\mathfrak{g}$-modules $V_{\lambda+\mu} \hookrightarrow V_\lambda \otimes V_\mu$, $v_{\lambda+\mu} \mapsto v_\lambda \otimes v_\mu$. Therefore we have dual surjective maps $V^*_\lambda \otimes V^*_\mu \to V^*_{\lambda+\mu}$, which define an algebra

$$\bar{Q}(d_1,\dots,d_s) = \bigoplus_{\lambda = m_1\omega_{d_1} + \dots + m_s\omega_{d_s}} V^*_\lambda.$$

Then one has an isomorphism of algebras $\bar{Q}(d_1, \dots, d_s) \simeq Q(d_1, \dots, d_s)$.

One can also embed the ring $Q(d_1, \dots, d_s)$ into a polynomial ring (thus proving that $I(d_1, \dots, d_s)$ is prime). Namely, let $Z_{i,j}$, $1 \le i, j \le n$ be a set of variables. Then the map

$$X_{j_1,\dots,j_d} \mapsto \det(Z_{l,j_l})_{1 \le l \le d} \tag{1.2}$$

induces the embedding $Q(d_1, \dots, d_s) \hookrightarrow \mathbb{C}[Z_{i,j}]_{1 \le i,j \le n}$.

The ring $Q(d_1, \dots, d_s)$ has a basis described in terms of the semistandard tableaux. Namely, for a partition $\lambda = (\lambda_1 \ge \dots \ge \lambda_{n-1} \ge 0)$ we denote by Y_λ the corresponding Young diagram. For example, $Y_{(7,5,4,2,2)}$ is the following diagram:



We number the rows and columns of Y_λ from up to down and from left to right. Thus, the boxes of Y_λ are labeled by pairs (i, j). We denote by μ_j the length of the j-th column.

A tableau T of shape λ is a filling of Y_λ with numbers $T_{i,j} \in \{1, \dots, n\}$. The number $T_{i,j}$ is attached to the box in the i-th row and j-th column. The numbers $T_{i,j}$ are subject to the condition: $i_1 < i_2$ implies $T_{i_1,j} < T_{i_2,j}$. A tableau is called semistandard if in addition $j_1 < j_2$ implies $T_{i,j_1} \le T_{i,j_2}$.

Let $P^+(d_1, \dots, d_s) = \mathbb{Z}_+\omega_{d_1} \oplus \dots \oplus \mathbb{Z}_+\omega_{d_s}$. Then we have a natural decomposition

$$Q(d_1, \dots, d_s) = \bigoplus_{\lambda \in P^+(d_1,\dots,d_s)} Q_\lambda(d_1, \dots, d_s) = \bigoplus_{\lambda \in P^+(d_1,\dots,d_s)} V_\lambda^*.$$

Recall that to a weight $\lambda = \sum_{i=1}^{n-1} m_i\omega_i$ one can attach a partition

$$(m_1 + \dots + m_{n-1}, m_2 + \dots + m_{n-1}, \dots, m_{n-1}),$$

which we denote by the same symbol λ.

Finally, for a tableau T of shape λ we define an element

$$X_T = \prod_{j=1}^{\lambda_1} X_{T_{1,j},\dots,T_{\mu_j,j}}$$

(note that $\lambda \in P^+(d_1, \dots, d_s)$ implies $\mu_j \in \{d_1, \dots, d_s\}$ for all j). Then the elements X_T labeled by the shape λ semistandard tableaux form a basis of $Q(d_1, \dots, d_s)$.

2. Abelian degenerations

We first recall the definition of the PBW-filtration on a highest weight module V_λ.

The space V_λ is equipped with the increasing PBW filtration F_s defined as follows:

$$F_0 = \mathbb{C}v_\lambda,\ F_s = \mathrm{span}\{x_1 \dots x_k v_\lambda :\ k \le s, x_i \in \mathfrak{n}^-\}.$$

In other words, $F_{s+1} = F_s + \mathfrak{n}^- F_s$. We denote by V_λ^a the associated graded space,

$$V_\lambda^a = F_0 \oplus \bigoplus_{s=1}^{\infty} F_s/F_{s-1}. \tag{2.1}$$

The superscript a stands for the abelian, see the explanation below. In what follows we write $V_\lambda^a = \bigoplus_{s\ge 0} V_\lambda^a(s)$, $V_\lambda^a(s) = F_s/F_{s-1}$. An element $x \in V_\lambda^a(s)$ is said to be homogeneous of the PBW degree s.

Let us describe the Lie algebra and Lie group acting on V_λ^a.

Definition 2.1. The Lie algebra $\mathfrak{g}^a$ is isomorphic to $\mathfrak{g}$ as a vector space and the bracket $[\cdot,\cdot]^a$ is given by the formulas

$$[h, f_\alpha]^a = -\alpha(h) f_\alpha,\ [h, e_\alpha]^a = \alpha(h) e_\alpha,\ [h, h_1]^a = 0 \qquad \forall h, h_1 \in \mathfrak{h}, \alpha \in Q_+,$$
$$[f_\alpha, f_\beta]^a = 0,\ [e_\alpha, e_\beta]^a = [e_\alpha, e_\beta] \qquad \forall \alpha, \beta \in Q_+,$$
$$[e_\alpha, f_\beta]^a = \begin{cases} [e_\alpha, f_\beta], \text{ if } \beta - \alpha \in Q_+, \\ 0, \text{ otherwise.} \end{cases}$$

It is straightforward to check that the bracket $[\cdot,\cdot]^a$ satisfies the Jacobi identity. The superscript a stays for the abelian, since the subalgebra $\mathfrak{n}^-$ is abelian with respect to the bracket $[\cdot,\cdot]^a$. In what follows we omit the superscript a in the bracket $[\cdot,\cdot]^a$ if it is clear what algebra $\mathfrak{g}$ or $\mathfrak{g}^a$ is considered.

Remark 2.2. The Lie algebra $\mathfrak{g}^a$ is a semidirect sum of the Borel subalgebra $\mathfrak{b}$ and an abelian ideal $(\mathfrak{n}^-)^a$, which is isomorphic to $\mathfrak{n}^-$ as a vector space. The action of $\mathfrak{b}$ on $(\mathfrak{n}^-)^a$ can be described as follows. The Lie algebra $\mathfrak{g}$ is naturally equipped with a structure of adjoint $\mathfrak{b}$-module and $\mathfrak{b} \hookrightarrow \mathfrak{g}$ is a submodule. Therefore we obtain a structure of quotient $\mathfrak{b}$-module on the space $(\mathfrak{n}^-)^a \simeq \mathfrak{g}/\mathfrak{b}$.

Remark 2.3. The Lie algebra $\mathfrak{g}^a$ can be considered as a degeneration of $\mathfrak{g}$. In fact, let $c^{++}_{\alpha,\beta}$, $c^{--}_{\alpha,\beta}$ and $c^{+-}_{\alpha,\beta}$ be the structure constants of $\mathfrak{g}$, i.e.

$$[e_\alpha, e_\beta] = c^{++}_{\alpha,\beta} e_{\alpha+\beta},\ [f_\alpha, f_\beta] = c^{--}_{\alpha,\beta} f_{\alpha+\beta},$$
$$[e_\alpha, f_\beta] = c^{+-}_{\alpha,\beta} f_{\beta-\alpha},\ \beta - \alpha \in Q_+,\ [e_\alpha, f_\beta] = c^{+-}_{\alpha,\beta} e_{\alpha-\beta},\ \alpha - \beta \in Q_+,$$
$$[e_\alpha, f_\alpha] = h_\alpha \in \mathfrak{h}.$$

Let $\mathfrak{g}(\varepsilon)$ be the subalgebra of $\mathfrak{g}$ spanned by the elements $e_\alpha(\varepsilon) = e_\alpha$, $f_\alpha(\varepsilon) = \varepsilon f_\alpha$, $\alpha \in Q_+$ and $\mathfrak{h}$. Obviously, $\mathfrak{g}(\varepsilon) = \mathfrak{g}$ for any $\varepsilon \neq 0$. Then one has

$$[e_\alpha(\varepsilon), e_\beta(\varepsilon)] = c^{++}_{\alpha,\beta} e_{\alpha+\beta}(\varepsilon),\ [f_\alpha(\varepsilon), f_\beta(\varepsilon)] = \varepsilon c^{--}_{\alpha,\beta} f_{\alpha+\beta}(\varepsilon),$$
$$[e_\alpha(\varepsilon), f_\beta(\varepsilon)] = c^{+-}_{\alpha,\beta} f_{\beta-\alpha}(\varepsilon),\ \beta - \alpha \in Q_+,$$
$$[e_\alpha(\varepsilon), f_\beta(\varepsilon)] = \varepsilon c^{+-}_{\alpha,\beta} e_{\alpha-\beta}(\varepsilon),\ \alpha - \beta \in Q_+,$$
$$[e_\alpha(\varepsilon), f_\alpha(\varepsilon)] = \varepsilon h_\alpha.$$

In the limit $\varepsilon \to 0$, the commutation relations above give the relations from Definition 2.1.

We now define the Lie group G^a, which corresponds to $\mathfrak{g}^a$. Let $M = \dim \mathfrak{n}$ and let $\mathbb{G}_a$ be the additive group of the field $\mathbb{C}$. The Lie group $\mathbb{G}_a^M$ can be naturally identified with the Lie group of the abelian Lie algebra $(\mathfrak{n}^-)^a \hookrightarrow \mathfrak{g}^a$. We note that B acts on $\mathfrak{g}$ via the restiction of the adjoint G-action. Since $\mathfrak{b} \hookrightarrow \mathfrak{g}$ is B-invariant, we obtain a structure of B-module on the vector space $(\mathfrak{n}^-)^a \simeq \mathfrak{g}/\mathfrak{b}$.

Definition 2.4. The Lie group G^a is a semidirect product $\mathbb{G}_a^M \rtimes B$ of the normal subgroup $\mathbb{G}_a^M$ and the Borel subgroup B. The action by conjugation of B on $\mathbb{G}_a^M$ is induced from the B-action on $(\mathfrak{n}^-)^a$ as above.

The following proposition is simple, but of the key importance for us. Recall the graded modules $V_\lambda^a = F_0 \oplus \bigoplus_{s\geq 1} F_s/F_{s-1}$ (see (2.1)).

Proposition 2.5. *For any $\lambda \in P_+$ the structure of $\mathfrak{g}$-module on V_λ induces the structures of $\mathfrak{g}^a$- and G^a-modules on V_λ^a.*

Proof. The action of the Lie algebra $\mathfrak{n}^-$ on V_λ induces the action of the abelian algebra $(\mathfrak{n}^-)^a \hookrightarrow \mathfrak{g}^a$ on V_λ^a. In fact, for any $\alpha \in Q_+$ we have $f_\alpha F_s \hookrightarrow F_{s+1}$. Therefore, we have the action of operators f_α on the graded space V_λ^a, mapping F_s/F_{s-1} to F_{s+1}/F_s. From definition of F_s we conclude that all such operators commute. Now consider the action of $\mathfrak{b}$ on V_λ. Since $xF_s \hookrightarrow F_s$ for any $x \in \mathfrak{b}$, we obtain the action of $\mathfrak{b}$ on the graded space V_λ^a mapping each quotient F_s/F_{s-1} to itself. Now it is easy to see that the action of abelian $(\mathfrak{n}^-)^a$ and of $\mathfrak{b}$ satisfy the relations from Definition 2.1. Thus we obtain a structure of $\mathfrak{g}^a$-module on V_λ^a. The group G^a acts on V_λ^a by the exponents of the operators from $\mathfrak{g}^a$. □

Remark 2.6. The space V_λ^a is a cyclic module over the universal enveloping algebra of $(\mathfrak{n}^-)^a \hookrightarrow \mathfrak{g}^a$ and therefore can be identified with a quotient of the polynomial algebra in variables f_α, $\alpha \in Q_+$ modulo some ideal $I(\lambda)$: $V_\lambda^a \simeq \mathbb{C}[f_\alpha]_{\alpha \in Q_+}/I(\lambda)$. The operators $e_\alpha \in \mathfrak{n}$ act on this quotient as differential operators. This ideal in type A was computed in [FFoL].

Example 2.7. Let $\mathfrak{g} = \mathfrak{sl}_n$, $\lambda = \omega_d$, $1 \leq d \leq n-1$. Recall the basis v_J of the module $V_{\omega_d} = \Lambda^d(\mathbb{C}^n)$. For a sequence $J = (j_1, \dots, j_d)$ we define the PBW degree of J by the formula

$$\deg J = \#\{r :\ j_r > d\}. \tag{2.2}$$

Then the s-th space of the PBW filtration F_s is spanned by the vectors v_J with $\deg J \le s$. Therefore, the images of the elements v_J with $\deg J = s$ define a basis of $F_s/F_{s-1} \hookrightarrow V^a_{\omega_d}$. We denote these images by the symbol v^a_J and the coordinates (dual basis) by X^a_J.

The action of $\mathfrak{g}^a$ on $V^a_{\omega_d}$ can be written explicitly. Namely, for $1 \le i \le j < n$ let $\alpha_{i,j} = \alpha_i + \dots + \alpha_j$ be positive roots of $\mathfrak{sl}_n$ and let $f_{i,j} = f_{\alpha_{i,j}}$, $e_{i,j} = e_{\alpha_{i,j}}$. Then it is easy to see that

$$f_{i,j} v^a_J = \begin{cases} \sum_{r=1}^d \delta_{i,j_r} v^a_{j_1,\dots,j_{r-1},j+1,j_{r+1},\dots,j_d}, & \text{if } i \le d, j \ge d, \\ 0, & \text{otherwise ;} \end{cases} \tag{2.3}$$

$$e_{i,j} v^a_J = \begin{cases} \sum_{r=1}^d \delta_{j+1,j_r} v^a_{j_1,\dots,j_{r-1},i,j_{r+1},\dots,j_d}, & \text{if } d < i \le j \text{ or } i \le j < d, \\ 0, & \text{otherwise .} \end{cases}$$

We note that $V^a_{\omega_d}$ is no longer isomorphic to $\Lambda^d(V^a_{\omega_1})$.

The representations V^a_λ for $\mathfrak{g} = \mathfrak{sl}_n$ were studied in [FFoL]. In particular, the following lemma is proved in [FFoL] (see the proof of Theorem 3.11):

Lemma 2.8. *Let $\lambda, \mu \in P_+$. Then*

$$V^a_{\lambda+\mu} \simeq \mathrm{U}(\mathfrak{g}^a)(v_\lambda \otimes v_\mu) \hookrightarrow V^a_\lambda \otimes V^a_\mu$$

as $\mathfrak{g}^a$-modules.

We now define the degenerate flag varieties $\mathcal{F}^a_\lambda$. Let $[v_\lambda] \in \mathbb{P}(V^a_\lambda)$ be the line $\mathbb{C}v_\lambda$.

Definition 2.9. The variety $\mathcal{F}^a_\lambda \hookrightarrow \mathbb{P}(V^a_\lambda)$ is a closure of the G^a-orbit of $[v_\lambda]$,

$$\mathcal{F}^a_\lambda = \overline{G^a[v_\lambda]} = \overline{\mathbb{G}^M v_\lambda} \hookrightarrow \mathbb{P}(V^a_\lambda).$$

We note that the orbit $G[v_\lambda] \hookrightarrow \mathbb{P}(V_\lambda)$ coincides with its closure, but the orbit $G^a[v_\lambda]$ does not. More precisely, since the Borel subgroup B stabilizes $[v_\lambda]$, the orbit $G^a[v_\lambda] = \mathbb{G}^M_a[v_\lambda]$ is an affine space $\mathbb{C}^{M_\lambda}$, where

$$M_\lambda = \#\{\alpha \in Q_+ : \ (\lambda, \alpha) > 0\}.$$

In fact,

$$\mathbb{G}^M_a[v_\lambda] = \left\{ \exp\left(\sum_{\alpha \in Q_+} c_\alpha f_\alpha \right) [v_\lambda], \ c_\alpha \in \mathbb{C} \right\}.$$

Since $f_\alpha v_\lambda = 0$ iff $(\lambda, \alpha) = 0$, we obtain $\mathbb{G}^M_a[v_\lambda] \simeq \mathbb{C}^{M_\lambda}$.

Remark 2.10. The variety $\mathcal{F}^a_\lambda$ is a $\mathbb{G}^{M_\lambda}_a$-equivariant compactification of the affine space $\mathbb{C}^{M_\lambda}$. In other words, $\mathcal{F}^a_\lambda$ are the so-called $\mathbb{G}^M_a$-varieties (see [HT], [A]).

3. PLÜCKER RELATIONS, COORDINATE RINGS AND PBW-TABLEAUX: THE DEGENERATE CASE

In this section we state main results of this paper. The proofs are given in the following sections.

Starting from this section we assume that $\mathfrak{g} = \mathfrak{sl}_n$. We first consider the varieties $\mathcal{F}^a_{\omega_d}$.

Lemma 3.1. *$\mathcal{F}^a_{k\omega_d} \simeq \mathcal{F}_{k\omega_d}$ for all $d = 1, \dots, n-1$, $k \geq 0$.*

Proof. We note that for a fixed d the elements f_α with $(\alpha, \omega_d) \neq 0$ pairwise commute in $\mathfrak{sl}_n$. Therefore,

$$\mathcal{F}_{\omega_d} = \overline{\left\{ \exp\left(\sum_{\alpha:(\alpha,\omega_d)\neq 0} c_\alpha f_\alpha \right) [v_{\omega_d}], c_\alpha \in \mathbb{C} \right\}} \simeq \mathcal{F}^a_{\omega_d}.$$

□

Remark 3.2. The reason that Lemma 3.1 holds for all fundamental weights is that all ω_d are cominuscule in type A (the radical, corresponding to each ω_d is abelian, see [FFL], [FL]). This is not true in general for other types, so for general $\mathfrak{g}$ the varieties $\mathcal{F}_{\omega_d}$ and $\mathcal{F}^a_{\omega_d}$ do not coincide for all d.

Because of Lemma 2.8, we have the following realization of the varieties $\mathcal{F}^a_\lambda$.

Proposition 3.3. *For $\lambda \in P_+$ let $1 \leq d_1 < \cdots < d_s < n$ be the set of all numbers d such that $(\lambda, \alpha_d) \neq 0$. Then*

$$\mathcal{F}^a_\lambda \simeq \overline{\mathbb{G}_a^M([v_{\omega_{d_1}}] \times \cdots \times [v_{\omega_{d_s}}])} \hookrightarrow Gr(d_1, n) \times \cdots \times Gr(d_s, n).$$

Remark 3.4. We note that since all f_α commute in V^a_λ, we have

$$\exp\left(\sum_{\alpha\in Q_+} c_\alpha f_\alpha \right) \left([v_{\omega_{d_1}}] \times \cdots \times [v_{\omega_{d_s}}] \right) = \tag{3.1}$$

$$\exp\left(\sum_{\alpha:(\alpha,\omega_{d_1})\neq 0} c_\alpha f_\alpha \right) [v_{\omega_{d_1}}] \times \cdots \times \exp\left(\sum_{\alpha:(\alpha,\omega_{d_s})\neq 0} c_\alpha f_\alpha \right) [v_{\omega_{d_s}}].$$

This formula is specific for the degenerate situation and is not true in the classical case. The reason is that the operators f_α with $(\alpha, \lambda) = 0$ act trivially on V^a_λ, but not on V_λ itself.

3.1. **Degenerate Plücker relations.**

We introduce the notation

$$\mathcal{F}^a(d_1, \dots, d_s) = \mathcal{F}^a_{\omega_{d_1}+\cdots+\omega_{d_s}}, \ 1 \leq d_1 < \cdots < d_s < n.$$

Definition 3.5. Let $I^a(d_1, \dots, d_s)$ be an ideal in the polynomial ring in variables $X^a_{j_1,\dots,j_d}$, $d = d_1, \dots, d_s$, $1 \leq j_1 < \cdots < j_d < n$, generated by the elements $R^{k;a}_{L,J}$ given below. These elements are labeled by a pair of numbers

$p \geq q$, $p, q \in \{d_1, \dots, d_s\}$, by an integer k, $1 \leq k \leq q$ and by sequences $L = (l_1, \dots, l_p)$, $J = (j_1, \dots, j_q)$, which are arbitrary subsets of the set $\{1, \dots, n\}$. The generating elements are given by the formulas

$$R^{k;a}_{L,J} = X^a_{l_1,\dots,l_p} X^a_{j_1,\dots,j_q} - \sum_{1 \leq r_1 < \dots < r_k \leq p} X^a_{l'_1,\dots,l'_p} X^a_{j'_1,\dots,j'_q}, \tag{3.2}$$

where the terms of $R^{k;a}_{L,J}$ are the terms of $R^k_{L,J}$ (1.1) (with a superscript a, to be precise) such that

$$\{l_{r_1}, \dots, l_{r_k}\} \cap \{q+1, \dots, p\} = \emptyset. \tag{3.3}$$

Remark 3.6. The initial term $X^a_{l_1,\dots,l_p} X^a_{j_1,\dots,j_q}$ is also subject to the condition (3.3), i.e. it is not present in $R^{k;a}_{L,J}$ if $\{j_1, \dots, j_k\} \cap \{q+1, \dots, p\} \neq \emptyset$.

Remark 3.7. Let us write the relation $R^k_{L,J}$ as $\sum_i X_{L^{(i)}} X_{J^{(i)}}$. Then $R^{k;a}_{L,J}$ is equal to the sum of terms $X^a_{L^{(i_0)}} X^a_{J^{(i_0)}}$ such that the sum of PBW degrees (2.2) $\deg L^{(i_0)} + \deg J^{(i_0)}$ is minimal among all sums $\deg L^{(i)} + \deg J^{(i)}$. In fact, first note that among the numbers $l_1, \dots, l_p$ there exists at least one k-tuple $l_{r_1}, \dots, l_{r_k}$, $r_1 < \dots r_k$ such that

$$\{l_{r_1}, \dots, l_{r_k}\} \cap \{q+1, \dots, p\} = \emptyset.$$

Hence without loss of generality we can assume that

$$\{j_1, \dots, j_k\} \cap \{q+1, \dots, p\} = \emptyset.$$

Now assume that a pair (L', J') is obtained from (L, J) by interchanging $(l_{r_1}, \dots, l_{r_k})$ with $(j_1, \dots, j_k)$. Then

$$\deg L' + \deg J' = \deg L + \deg J + \#\{i : \ q+1 \leq l_{r_i} \leq p\}.$$

Therefore, to obtain $R^{k;a}_{L,J}$ from $R^k_{L,J}$, one has to pick the terms of minimal PBW-degree.

Remark 3.7 implies the following:

Corollary 3.8. *The ideal $I^a(d_1, \dots, d_s)$ is homogeneous with respect to the PBW-degree.*

We now work out several examples.

Example 3.9. Let $s = 1$. Then $I^a(d) = I(d)$, since there are no numbers l such that $d+1 \leq l \leq d$ and thus $R^{k;a}_{L,J} = R^k_{L,J}$ (up to a superscript a in the notations of variables X_J).

Example 3.10. Let $\mathfrak{g} = \mathfrak{sl}_3$. Then the only nontrivial ideal is $I^a(1,2)$. It is generated by a single element

$$R^{1;a}_{(1,2),(3)} = X^a_{1,2} X^a_3 + X^a_{2,3} X^a_1.$$

We note that

$$R^1_{(1,2),(3)} = X_{1,2} X_3 - X_{1,3} X_2 + X_{2,3} X_1.$$

The middle term is missing in $R^{1;a}_{(1,2),(3)}$, because

$$\deg(1,2)+\deg(3)=\deg(2,3)+\deg(1)=1, \text{ but } \deg(1,3)+\deg(2)=2,$$

or, equivalently, because 2 satisfies $1+1\le 2\le 2$ (here $q=1, p=2$).

Example 3.11. Let $\mathfrak{g}=\mathfrak{sl}_4$. Consider the ideal $I^a(1,2)$. This ideal is generated by the following elements:

$$R^{1;a}_{(1,2),(3)}=X^a_{1,2}X^a_3+X^a_{2,3}X^a_1,\ R^{1;a}_{(1,2),(4)}=X^a_{1,2}X^a_4+X^a_{2,4}X^a_1,$$
$$R^{1;a}_{(1,3),(4)}=X^a_{1,3}X^a_4-X^a_{1,4}X^a_3+X^a_{3,4}X^a_1,\ R^{1;a}_{(2,3),(4)}=X^a_{2,3}X^a_4-X^a_{2,4}X^a_3,$$
$$R^{1;a}_{(1,2),(3,4)}=X^a_{1,2}X^a_{3,4}-X^a_{1,3}X^a_{2,4}+X^a_{2,3}X^a_{1,4}.$$

We note that the relation $R^{1;a}_{(1,2),(3,4)}$ is exactly the one defining the Grassmanian $Gr(2,4)$.

Example 3.12. For an arbitrary n, consider the ideal $I^a(1,n-1)$. This ideal is generated by a single element

$$R^{1;a}_{(1,\dots,n-1),(n)}=X^a_{1,\dots,n-1}X^a_n-(-1)^nX^a_{2,\dots,n}X^a_1.$$

In the classical case the corresponding relation is given by

$$R^{1}_{(1,\dots,n-1),(n)}=\sum_{i=1}^n(-1)^{n-i}X_{1,\dots,\hat{i}\dots,n}X_i.$$

The following theorem is the main statement of our paper, it will be proved in Section 5.

Theorem 3.13. *The variety* $\mathcal{F}^a(d_1,\dots,d_s)\hookrightarrow\times_{i=1}^s\mathbb{P}(\Lambda^{d_i}\mathbb{C}^n)$ *is defined by the ideal* $I^a(d_1,\dots,d_s)$.

Example 3.14. Let $s=2$, $d_1=1$, $d_2=n-1$. The variety $\mathcal{F}^a(1,n-1)$ is embedded into the product $\mathbb{P}^{n-1}\times\mathbb{P}^{n-1}=\mathbb{P}(V^a_{\omega_1})\times\mathbb{P}(V^a_{\omega_{n-1}})$. The elements f_α acting nontrivially on $V^a_{\omega_1}$ are $f_{1,i}$, $i=1,\dots,n-1$, they act as $f_{1,i}v^a_j=\delta_{1,j}v^a_{i+1}$. Similarly, the elements f_α acting nontrivially on $V_{\omega_{n-1}}$ are $f_{i,n-1}$, $i=1,\dots,n-1$, they act as

$$f_{i,n-1}v^a_{1,\dots,\hat{j}\dots,n}=(-1)^{n-i+1}\delta_{j,n}v^a_{1,\dots,\hat{i}\dots,n}.$$

Therefore the coordinates of a point

$$\exp\left(\sum_{i=1}^{n-1}c_{1,i}f_{1,i}+\sum_{i=1}^{n-1}c_{i,n-1}f_{i,n-1}\right)\left([v_{\omega_1}]\otimes[v_{\omega_{n-1}}]\right)$$

are given by

$$X^a_1=1, X^a_i=c_{1,i-1}\ (i>1),\ X^a_{1,\dots,n-1}=1, X^a_{1,\dots,\hat{i}\dots,n}=(-1)^{n-i+1}c_{i,n-1}.$$

Thus, the open orbit of $\mathcal{F}^a(1, n-1)$ is the following set of points of the product of two projective spaces:

$$\left[v_1^a + \sum_{i=2}^{n} c_{1,i-1} v_i^a\right] \times \left[v_{1,\dots,n-1}^a + \sum_{i=1}^{n-1} (-1)^{n-i-1} c_{i,n-1} v_{1,\dots\hat{i},\dots,n}^a\right].$$

Therefore, the closure of the open cell is defined by a single algebraic relation

$$X_{1,\dots,n-1}^a X_n^a - (-1)^n X_{2,\dots,n}^a X_1^a = 0,$$

which agrees with Example 3.12. Recall that the classical variety $\mathcal{F}(1, n-1)$ is defined by a single relation $R^1_{(1,\dots,n-1),(n)}$.

It turns out that our degeneration is flat. We will prove the following proposition.

Proposition 3.15. *The varieties $\mathcal{F} = \mathcal{F}(d_1, \dots, d_s)$ and $\mathcal{F}^a = \mathcal{F}^a(d_1, \dots, d_s)$ can be connected by a flat family $\mathcal{F}^t$ such that $\mathcal{F}^0 \simeq \mathcal{F}^a$ and $\mathcal{F}^t \simeq \mathcal{F}$, if $t \neq 0$.*

3.2. **Coordinate rings: quotient polynomial rings.** Define a quotient algebra

$$Q^a(d_1, \dots, d_s) = \mathbb{C}[X_{i_1,\dots,i_d}^a]/I^a(d_1, \dots, d_s),$$

where the variables $X_{i_1,\dots,i_d}^a$ are labeled by sequences $1 \le i_1 < \dots < i_d \le n$, $d = d_1, \dots, d_s$.

Theorem 3.16. *The algebra $Q^a(d_1, \dots, d_s)$ is isomorphic to the algebra*

$$\bigoplus_{\lambda \in P^+(d_1,\dots,d_s)} (V_\lambda^a)^*,$$

where the multiplication $(V_\lambda^a)^* \otimes (V_\mu^a)^* \to (V_{\lambda+\mu}^a)^*$ is induced by the embedding $V_{\lambda+\mu}^a \hookrightarrow V_\lambda^a \otimes V_\mu^a$ (see Lemma 2.8).

3.3. **PBW-semistandard tableaux: polynomial subring.** A crucial statement about the ideal I^a is that it is prime. As in the classical case this follows from the realization of the quotient algebra as a subalgebra of the polynomial algebra. In the classical case the key role in the whole picture plays a notion of a semistandard tableau. We introduce the PBW-version here.

Recall that for a partition $\lambda = (\lambda_1 \ge \dots \ge \lambda_{n-1} \ge 0)$ we denote by Y_λ the corresponding Young diagram. We denote by μ_j the length of the j-th column.

Definition 3.17. A PBW-tableau of shape λ is a filling $T_{i,j}$ of the Young diagram Y_λ with numbers $1, \dots, n$. The number $T_{i,j} \in \{1, \dots .n\}$ is attached to the box (i, j). The filling $T_{i,j}$ has to satisfy the following properties:

- If $T_{i,j} \le \mu_j$, then $T_{i,j} = i$,
- If $i_1 < i_2$ and $T_{i_1,j} \neq i_1$, then $T_{i_1,j} > T_{i_2,j}$.

Remark 3.18. Usually the entries of a tableau increase from up to down and from left to right. The reason why it is more natural for us to change the directions is two fold: first, we want to make difference between numbers, which are smaller than the length of a column and the others. Second, our convention better fits to the formalism of Vinberg's patterns [V], [FFoL], see the proof of Proposition (4.5).

Remark 3.19. Let us explain the reason why we want $T_{i,j} \le \mu_j$ to imply $T_{i,j} = i$. Assume that Y_λ consists of one column of length d. Then we have $V_\lambda^a \simeq V_{\omega_d}^a$. As in the classical case we want the shape λ PBW-tableaux to label the standard basis of $V_{\omega_d}^a$. So let us take a basis vector $v_J^a = v_{j_1,\dots,j_d}^a$, $j_1 < \cdots < j_d$. This vector is obtained from the highest weight vector $v_{1,\dots,d}^a$ (which is the image of $v_1 \wedge \cdots \wedge v_d$) by application of several operators f_α. Note that the operators $f_{i,j}$ with $1 \le i \le j < d$ act trivially on $V_{\omega_d}^a$. Therefore, whenever a number $j \le d$ appears in J, it means that no f_α has been applied to v_j. Hence it is natural to reorder the vectors v_j in the product $v_{j_1} \wedge \cdots \wedge v_{j_d}$ in such a way that if $j_r \le d$ then j_r appears at the j_r-th place.

We note that $T_{i,j} \ge i$ for any pair i, j.

Definition 3.20. A semistandard PBW-tableau of shape λ is a PBW-tableau of shape λ subject to the condition:

- for any $j > 1$ and any i there exists $i_1 \ge i$ such that $T_{i_1,j-1} \ge T_{i,j}$.

We denote the set of semistandard PBW-tableaux of shape λ by ST_λ^a.

The shape λ semistandard PBW-tableaux parametrize a basis of V_λ^a. In particular

Proposition 3.21. *The number of semistandard PBW-tableaux is equal to* $\dim V_\lambda$.

Example 3.22. Let $\lambda = (\underbrace{1,\dots,1}_{d})$. Then each sequence $1 \le i_1 < \cdots < i_d \le n$ gives rise to a semistandard PBW-tableau of shape (d).

Example 3.23. Let $\lambda = (k)$ be a one part partition. Then the shape λ semistandard PBW-tableaux are of the following form:

$$\begin{array}{|c|c|c|c|c|c|c|c|c|c|}\hline n & \cdots & n & \quad\cdots\quad & 2 & \cdots & 2 & 1 & \cdots & 1 \\ \hline\end{array}$$

Here a number $i = 1, \dots, n$ appears a_i times, $a_i \ge 0$ and $\sum_{i=1}^n a_i = k$. In particular, the number of semistandard PBW-tableaux is equal to $\dim S^k(\mathbb{C}^n) = \dim V_{k\omega_1}$.

Example 3.24. Let $\mathfrak{g} = \mathfrak{sl}_3$, $\lambda = (2,1)$. Then the list of the semistandard PBW-tableaux is as follows:

$$\begin{array}{|c|c|}\hline 1 & 1 \\ \hline 2 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 1 & 2 \\ \hline 2 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 1 & 1 \\ \hline 3 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 1 & 2 \\ \hline 3 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 1 & 3 \\ \hline 3 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 3 & 1 \\ \hline 2 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 3 & 2 \\ \hline 2 \\ \cline{1-1}\end{array}\ ,\ \begin{array}{|c|c|}\hline 3 & 3 \\ \hline 2 \\ \cline{1-1}\end{array}\ .$$

Example 3.25. Let $\mathfrak{g} = \mathfrak{sl}_3$ and $\lambda = (m_1 + m_2, m_2)$ be an arbitrary partition. Then the semistandard PBW-tableaux are of the form

$$\begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|}\hline 1 & \dots & 1 & 3 & \dots & 3 & 1 & \dots & 1 & 3 & \dots & 3 & 2 & \dots & 2 & 1 & \dots & 1 \\ \hline 3 & \dots & 3 & 2 & \multicolumn{4}{c|}{\dots\dots} & 2 & \multicolumn{9}{c}{} \\ \cline{1-9} \end{array},$$

with the restriction $T_{1,m_2+1} \neq 3$ unless $T_{1,m_2} = 3$. More precisely,

$$T_{1,1} = \dots = T_{1,k_1} = T_{1,k_1+k_2+1} = \dots = T_{1,m_2} = \\ T_{1,m_2+k_3+k_4+1} = \dots = T_{1,m_1+m_2} = 1,$$

$$T_{1,m_2+k_3+1} = \dots = T_{1,m_2+k_3+k_4} = T_{2,k_2+1} = \dots = T_{2,m_2} = 2,$$

$$T_{2,1} = \dots = T_{2,k_1} = T_{1,k_1+1} = \dots = T_{1,k_1+k_2} = \\ T_{1,m_2+1} = \dots = T_{m_2+1+k_3} = 3,$$

$k_1, k_2, k_3, k_4 \geq 0$ and $k_3 = 0$ unless $k_1 + k_2 = m_2$. One can easily show that the number of such tableaux is equal to $(m_1 + 1)(m_2 + 1)(m_1 + m_2 + 1)/2$, which coincides with the dimension of $V_{m_1\omega_1+m_2\omega_2}$.

Given a PBW-tableau T, we define an element D^a_T in the ring $\mathbb{C}[Z_{i,j}]$, $i < j$. First, assume that we have a sequence of numbers $1 \leq i_1 < \dots < i_d \leq n$. Let $m \leq d$ be a number such that $i_m \leq d$ and $i_{m+1} > d$. Then $D^a_{i_1,\dots,i_d}$ is the determinant of the matrix $D^a_{\alpha,\beta}$, $\alpha, \beta = 1, \dots, d$

$$D^a_{\alpha,\beta} = \begin{cases} Z_{\alpha,i_\beta}, \text{ if } \alpha \neq i_1, \dots, i_m, \beta > m, \\ 0, \text{ if } \alpha \neq i_1, \dots, i_m, \beta \leq m, \\ 1, \text{ if } \alpha = i_1, \dots, i_m, \beta = \alpha, \\ 0, \text{ if } \alpha = i_1, \dots, i_m, \beta \neq \alpha. \end{cases} \tag{3.4}$$

Now assume we are given a length d column of pairwise distinct numbers. Then there exists a permutation $\sigma \in S_d$ such that σT has increasing entries. We then define $D^a_T = (-1)^\sigma D^a_{\sigma T}$. If λ has more than one column, the element D^a_T is the product of of the polynomials, corresponding to its columns.

Proposition 3.26. *For any relation $R^{k;a}_{L,J}$ one has $R^{k;a}_{L,J}(D^a_{i_1,\dots,i_d}) = 0$.*

We will prove the following proposition:

Proposition 3.27. *Fix a partition λ. The polynomials D^a_T, $T \in ST^a_\lambda$, are linearly independent*

For a tableau T of shape λ we define a n element $X^a_T \in Q^a(d_1, \dots, d_s)$ by the formula

$$X^a_T = \prod_{j=1}^{\lambda_1} X^a_{T_{1,j}} \dots X^a_{T_{\mu_j,j}}.$$

Proposition 3.27 allows to prove the following theorem.

Theorem 3.28. *The elements X^a_T, $T \in ST^a_\lambda$, form a basis of $Q^a(d_1, \dots, d_s)$.*

Recall the notation

$$P^+(d_1,\dots,d_s)=\{m_1\omega_{d_1}+\dots+m_s\omega_{d_s},\ m_i\in\mathbb{Z}_{\ge 0}\}.$$

Corollary 3.29. $Q^a(d_1,\dots,d_s)\simeq\bigoplus_{\lambda\in P^+(d_1,\dots,d_s)}(V_\lambda^a)^*$.

4. Coordinate rings and PBW-tableaux

In this section we start proving the results from the previous section. Throughout the section we fix a sequence of integers $1\le d_1<\dots<d_s<n$ and omit this sequence in the notations writing Q^a, I^a etc instead of $Q^a(d_1,\dots,d_s)$, $I^a(d_1,\dots,d_s)$ etc.

Consider the spaces

$$\bar{Q}^a=\bigoplus_{\lambda\in P^+(d_1,\dots,d_s)}(V_\lambda^a)^*.$$

This space can be endowed with the structure of an algebra with the multiplication $(V_\lambda^a)^*\otimes(V_\mu^a)^*\to(V_{\lambda+\mu}^a)^*$ coming from the embedding $V_{\lambda+\mu}^a\hookrightarrow V_\lambda^a\otimes V_\mu^a$. We note that $\bar{Q}^a$ is PBW-graded, since the embeddings $V_{\lambda+\mu}^a\hookrightarrow V_\lambda^a\otimes V_\mu^a$ is compatible with the PBW-grading. The algebra $\bar{Q}^a$ is generated by the subspaces $(V_{\omega_{d_i}}^a)^*$. Recall the elements $X_{i_1,\dots,i_d}^a\in(V_{\omega_d}^a)^*$.

Lemma 4.1. *The elements $X_{i_1,\dots,i_{d_l}}^a$, $l=1,\dots,s$ satisfy the relations $R_{L,J}^{k;a}$ in $\bar{Q}^a$.*

Proof. The elements $X_{i_1,\dots,i_{d_l}}$, $l=1,\dots,s$ do satisfy relations $R_{L,J}^k$ in the classical algebra $Q=\bigoplus_{\lambda\in P^+}V_\lambda^*$ and the relations $R_{L,J}^{k;a}$ are simply the lowest degree terms with respect to the PBW grading (see Remark 3.7). Since the algebra $\bar{Q}^a$ is PBW graded, our lemma follows. $\square$

Recall the algebra $Q^a=\mathbb{C}[X_{i_1,\dots,i_d}^a]/I^a$, $d=d_1,\dots,d_s$, $1\le i_1<\dots<i_d\le n$.

Corollary 4.2. *There is a surjection of algebras $Q^a\to\bar{Q}^a$.*

We will prove that this surjection is an isomorphism. Consider the decomposition

$$Q^a=\bigoplus_{\lambda\in P^+(d_1,\dots,d_s)}Q_\lambda^a,$$

where for $\lambda=m_1\omega_{d_1}+\dots+m_s\omega_{d_s}$ the subspace Q_λ^a is spanned by the monomials $X_{J(1)}^a\dots X_{J(r)}^a$, where

$$\#\{i:\ J^{(i)}\text{ has }d_l\text{ entries}\}=m_l,\ l=1,\dots,s.$$

Our goal is to show that $\dim Q_\lambda^a=\dim V_\lambda$.

Recall definitions 3.17, 3.20 of the PBW-tableau and of the semistandard PBW-tableau.

Lemma 4.3. *Let T be a semistandard PBW-tableau. Then:*

- *If* $j_1 < j_2$ *and* $T_{i,j_1} = T_{i,j_2} \ne i$, *then* $T_{i,j} = T_{i,j_1}$ *for all* $j_1 \le j \le j_2$.
- *If* $i_1 \le i_2$, $j_1 \le j_2$ *and* $T_{i_1,j_1} \ne i_1$, $T_{i_2,j_2} \ne i_2$, *then* $T_{i_1,j_1} \ge T_{i_2,j_2}$. *If in addition* $i_1 < i_2$, *then* $T_{i_1,j_1} > T_{i_2,j_2}$.

Proof. Let us prove the first statement. Assume that there exists j, $j_1 < j < j_2$ such that $T_{i,j} \ne T_{i,j_1}$. Then $T_{i,j} = i$, because otherwise from the definition of a PBW-tableau we have $T_{i,j_1} \ge T_{i,j} \ge T_{i,j_2}$ and one of the inequalities is strict, which is a contradiction. So $T_{i,j} < T_{i,j_2}$. By definition of a semistandard PBW-tableau, there exists a sequence $i_1 \le i_2 \le \dots \le i_{j_2-j_1}$ such that $i_1 \ge i$ and

$$T_{i,j_2} \le T_{i_1,j_2-1} \le T_{i_2,j_2-2} \le \dots \le T_{i_{j_2-j_1},j_1}.$$

Since $T_{i,j} < T_{i,j_2}$, we have $i_{j_2-j} > i$ and hence $i_{j_2-j_1} > i$. This gives

$$T_{i,j_2} \le T_{i_{j_2-j_1},j_1} < T_{i,j_1},$$

which gives a contradiction.

The second statement of the lemma can be proved similarly. $\square$

Corollary 4.4. *Let* T *be a semistandard PBW-tableau. For* $l > k$ *let* $A_{col}(k,l)$ *be the minimal number such that* $T_{k,A_{col}(k,l)+1} = l$. *Then*

- *there exists* $i \ge k$ *such that* $T_{i,A_{col}(k,l)} \ge l$;
- *for* $j > A_{col}(k,l)$ *and* $i > k$ $T_{i,j} < l$.

Proposition 4.5. *The number of shape* λ *semistandard PBW-tableaux is equal to the dimension of* V_λ.

Proof. Let $\lambda = \sum_{i=1}^{n-1} m_i\omega_i$. Recall (see [FFoL], [V]) that the dimension of V_λ is equal to the number of Vinberg's configurations. A configuration is a set of numbers $(s_{i,j})_{1\le i\le j\le n-1} \in \mathbb{Z}_{\ge 0}$ or, equivalently, a set of numbers s_α labeled by positive roots α (we identify a root $\alpha = \alpha_i + \dots + \alpha_j$, $i \le j$, with the pair i, j). A configuration is said to be Vinberg's configuration, if the following condition holds: for any Dyck path $\mathbf{p} = (p_0, \dots, p_k)$ with $p(0) = \alpha_i$, $p(k) = \alpha_j$ we have

$$\sum_{l=0}^{k} s_{p_l} \le m_i + \dots + m_j. \tag{4.1}$$

Recall that a Dyck path $\mathbf{p} = (p_0, \dots, p_k)$ is a sequence of roots of $\mathfrak{sl}_n$,

$$p_l = \alpha_{i_l,j_l} = \alpha_{i_l} + \dots + \alpha_{j_l},\ i_l \le j_l,$$

such that either $(i_{l+1} = i_l,\ j_{l+1} = j_l + 1)$ or $(i_{l+1} = i_l + 1,\ j_{l+1} = j_l)$. Let S_λ be the set of Vinberg's configurations. We construct a one-to-one map ψ from S_λ to the set ST^a_λ of semistandard PBW-tableaux of shape λ.

Let $\mathbf{s} \in S_\lambda$ be some configuration. We define a tableau $\psi(\mathbf{s})$ in the following way. We fill the diagram Y_λ from bottom to top. Let us start with the $(n-1)$-st row (if there is no such row, the procedure described below is empty). We define

$$T_{n-1,1} = \dots = T_{n-1,s_{n-1,n-1}} = n$$

and fill the rest of the $(n-1)$-st row with the numbers $n-1$.

Now assume that the rows from $k+1$ to n are already filled. We fill the k-th row in the following way. Let $A_{col}(k,n)$ be the maximal number such that there exists $k \le i \le n-1$ such that $T_{i,A_{col}(k,n)} = n$ (since nothing is present in the k-th row at the moment, we can say $k < i$). We then define

$$T_{k,A_{col}(k,n)+1} = \dots = T_{k,A_{col}(k,n)+s_{k,n-1}} = n.$$

In general, we fill the k-th row with the numbers $n, n-1, \dots, k+1$ in the decreasing order in the following way. Assume that all numbers $n, \dots, l+1$ we want to put in the k-th row are already there. Let us explain how do we put the numbers l. Let $A_{col}(k,l)$ be the maximal number such that there exists $i = k, \dots, n$ such that $T_{i,A_{col}(k,l)} \ge l$. We then define

$$T_{k,A_{col}(k,l)+1} = \dots = T_{k,A_{col}(k,l)+s_{k,l-1}} = l.$$

After all iterations of the procedure above (corresponding to the numbers $l = n, \dots, k+1$) we fill the rest boxes of the k-th row with numbers k.

Note that the procedure above fills the k-th row of a tableau with numbers greater than or equal to k. We have to show that the construction above is well-defined and produces a semistandard PBW-tableau. For this we have to show that the numbers we fill the k-th row with fit into it. Since the length of the k-th row is equal to $m_k + \dots + m_{n-1}$, it suffices to show that for all $l > k$

$$A_{col}(k,l) + s_{k,l-1} \le m_k + \dots + m_{n-1}. \tag{4.2}$$

We define

$$A_{row}(k,l) = \min\{i : \ T_{i,A_{col}(k,l)} \ge l\}.$$

We set $A(k,l) = (A_{row}(k,l), A_{col}(k,l))$.

Assume first that $A_{row}(k,l) \ge l$. Then

$$A_{col}(k,l) \le m_l + \dots + m_{n-1}$$

and since $s_{k,l-1} \le m_k + \dots + m_{l-1}$, the inequality (4.2) holds. Now let $A_{row}(k,l) < l$. Then since $T_{A(k,l)} \ge l$, we have $T_{A(k,l)} \ne A_{row}(k,l)$. Set

$$k_1 = A_{col}(k,l), \quad l_1 = T_{A(k,l)}.$$

Note that $k_1 \ge k$ and $l_1 \ge l$. Then $A_{row}(k,l) = A_{row}(k_1,l_1) + s_{k_1,l_1-1}$. Thus (4.2) is equivalent to

$$A_{col}(k_1,l_1) + s_{k_1,l_1-1} + s_{k,l-1} \le m_k + \dots + m_{n-1}. \tag{4.3}$$

As on the previous step, assume first that $A_{row}(k_1,l_1) \ge l_1$. Then

$$A_{col}(k_1,l_1) \le m_{l_1} + \dots + m_{n-1}$$

and since $s_{k,l-1} + s_{k_1,l_1-1} \le m_k + \dots + m_{l_1-1}$ (recall that $k_1 \ge k$ and $l_1 \ge l$ and therefore the points $(k, l-1)$ and (k_1, l_1-1) can be connected by a Dyck path), the inequality (4.3) holds. So let $A_{row}(k_1,l_1) < l_1$. Define

$$k_2 = A_{col}(k_1,l_1), \quad l_2 = T_{A(k_1,l_1)}.$$

We note that $k_2 \geq k_1$ and $l_2 \geq l_1$. The (4.3) is equivalent to

$$A_{col}(k_2,l_2) + s_{k_2,l_2-1} + s_{k_1,l_1-1} + s_{k,l-1} \leq m_k + \dots + m_{n-1}. \tag{4.4}$$

Note that the points (k,l), (k_1,l_1) and (k_2,l_2), can be connected by a Dyck path.

We continue in the same way until we get $A_{col}(k_p,l_p) = 0$. The corresponding inequality (analogous to (4.2), (4.3) and (4.4)) follows from the inequality on the sum of $s_\bullet$ on a single Dyck path.

Now we need to construct an opposite map ψ^{-1} from ST^a_λ to S_λ. Given a tableau, we define

$$s_{k,l} = \#\{j : \ T_{k,j} = l+1\}. \tag{4.5}$$

Taking into account Corollary 4.4 we prove in the same way as above that $\mathbf{s}$ defined by (4.5) is an element of S_λ. Finally, we note that the two maps constructed above are inverse to each other. □

In order to establish the isomorphism $Q^a \simeq \bar{Q}^a$ and to prove that I^a is prime, we introduce one more algebra $\tilde{Q}^a$ (actually, we will show that $\tilde{Q}^a \simeq Q^a$). Let $\tilde{Q}^a$ be an algebra inside the polynomial ring $\mathbb{C}[Z_{i,j}]_{1\le i<j\le n}$ generated by the elements $D^a_{i_1,\dots,i_d}$ (see (3.4)).

Lemma 4.6. *For all k, L and J one has $R^{k;a}_{L,J}(D^a_{i_1,\dots,i_d}) = 0$ in $\mathbb{C}[Z_{i,j}]_{1\le i<j\le n}$.*

Proof. We note that the definition of $D^a_{i_1,\dots,i_d}$ can be given in the form of the equality in $V^a_{\omega_d}$:

$$\exp\left(\sum_{i<j} Z_{i,j} f_{i,j-1}\right)[v^a_{1,\dots,d}] = \sum_{i_1<\dots<i_d} D^a_{i_1,\dots,i_d} v^a_{i_1,\dots,i_d}.$$

The statement that these coefficients satisfy relations from I^a is proved in the following section, see Lemma 5.4. □

Corollary 4.7. *There exists a surjection of algebras $Q^a \to \tilde{Q}^a$.*

Lemma 4.8. *Fix a partition λ. Then the elements D^a_T, $T \in ST^a_\lambda$, are linearly independent.*

Proof. Let T be a semistandard PBW-tableau. Note that D^a_T contains a monomial

$$M^a_T = \prod_{i,j:\ T_{i,j}\neq i} Z_{i,T_{i,j}}.$$

It is convenient to rewrite this monomial in terms of the corresponding Vinberg's configuration $\mathbf{s} = \psi^{-1}(T)$ (see the bijection in the proof of Proposition 4.5). Namely it is easy to see that

$$M^a_{\mathbf{s}} = \prod_{i\le l} Z^{s_{i,l}}_{i,l+1}.$$

Let us consider the following order on the set S_λ. We say $\mathbf{s} > \mathbf{t}$ if there exists a pair i_0, l_0 such that $s_{i_0,l_0} > t_{i_0,l_0}$ and

$$s_{i,l} = t_{i,l} \text{ if } (i < i_0 \text{ or } i = i_0, l < l_0).$$

Note that if $\mathbf{s} > \mathbf{t}$, then the monomial $M^a_{\mathbf{s}}$ does not appear in $D^a_{\psi(\mathbf{t})}$. Therefore the elements D^a_T are linearly independent. □

Recall that for a shape λ tableau T we have the elements X^a_T defined by

$$X^a_T = \prod_{j=1}^{\lambda_1} X^a_{T_{1,j},\dots,T_{\mu_j,j}}.$$

Lemma 4.9. *The elements X^a_T, $T \in ST^a_\lambda$, span Q^a and hence $\tilde{Q}^a$ and $\bar{Q}^a$.*

Proof. We introduce an order on the set of tableaux of shape λ. We say $T^{(1)} > T^{(2)}$ if there exist i_0, j_0 such that $T^{(1)}_{i_0,j_0} > T^{(2)}_{i_0,j_0}$ and

$$T^{(1)}_{i,j} = T^{(2)}_{i,j} \text{ if } j > j_0 \text{ or } (j = j_0, i > i_0).$$

Assume that we are given a PBW-tableau T, which is not semistandard. Recall that the elements $D^a_{i_1,\dots,i_d}$ do satisfy relations from I^a. Using these relations we rewrite D^a_T in terms of smaller tableaux.

Assume that the condition from definition 3.20 is violated. Then there exist k, j such that $T_{k,j+1} > k$ and for all $i \geq k$ $T_{k,j+1} > T_{i,j}$. Let $p = \mu_j$, $q = \mu_{j+1}$. Since $T_{k,j+1} > T_{p,j} \geq p$, none of the elements $T_{1,j+1}, \dots, T_{k,j+1}$ belong to the set $\{q+1, \dots, p\}$ and hence the term

$$X^a_{T_{1,j},\dots,T_{k,j}} X^a_{T_{1,j+1},\dots,T_{k,j+1}}$$

is present in the relation

$$R^{k;a}_{(T_{1,j},\dots,T_{p,j}),(T_{1,j+1},\dots,T_{q,j+1})}. \tag{4.6}$$

Let us look at other terms in (4.6). Assume we pick arbitrary numbers $1 \leq i_1 < \dots < i_k \leq p$ and interchange

$$T_{i_1,j}, \dots, T_{i_k,j} \text{ with } T_{1,j+1}, \dots, T_{k,j+1}.$$

Since $T_{i,j} < T_{k,j+1}$ for all $i \geq k$, we have (after interchanging) at least $q - k + 1$ elements $T_{i_k,j}, T_{k+1,j+1}, \dots, T_{q,j+1}$ which are smaller than $T_{k,j+1}$. Therefore the lowest element in the column j_2, which differs from the one in T is smaller. Therefore, after interchanging the resulting tableau becomes smaller. □

Theorem 4.10. *We have isomorphisms $Q^a \simeq \bar{Q}^a \simeq \tilde{Q}^a$. The elements X^a_T labeled by semistandard PBW-tableaux T form a basis of Q^a.*

Proof. In fact, Lemma 4.9 and Proposition 4.5 give the inequalities $\dim Q^a_\lambda \leq \dim V_\lambda$ and $\dim \tilde{Q}^a_\lambda \leq \dim V_\lambda$. From Lemma 4.8 we obtain $\dim \tilde{Q}^a_\lambda = \dim V_\lambda$. Since $\dim \bar{Q}^a_\lambda = \dim V_\lambda$, Corollary 4.2 finishes the proof. □

Corollary 4.11. *The ideal I^a is prime.*

Proof. Follows from the isomorphism $Q^a \simeq \tilde{Q}^a$, since $\tilde{Q}^a$ is a subalgebra in the polynomial algebra. □

We close this section with the following proposition.

Proposition 4.12. *The elements X_T, $T \in ST^a_\lambda$, form a basis of Q.*

Proof. Since $\# ST^a_\lambda = \dim V_\lambda$, it suffices to prove that the elements X_T, $T \in ST^a_\lambda$, span Q. Recall that in the proof of Lemma 4.9 we have introduced an order on the set of tableau and expressed each X^a_T, $T \notin ST^a_\lambda$ as a linear combination of $X_{T'}$ with T' smaller than T. For this we have used the relations $R^{k;a}_{L,J}$. We note that

$$R^k_{L,J} = \tilde{R}^{k;a}_{L,J} + \sum_i X_{L^{(i)}} X_{J^{(i)}},$$

where $\deg J^{(i)} + \deg L^{(i)} > \deg J + \deg L$ and $\tilde{R}^{k;a}_{L,J}$ is obtained from $R^{k;a}_{L,J}$ simply by omitting superscripts a in variables X^a_J. Therefore in Q any element X_T, $T \notin ST^a_\lambda$ can be expressed as a linear combination of $X_{T'}$, where T' is smaller than T or the sum of PBW-degrees of columns of T' is bigger than that of T. Since the sum of PBW-degrees of fixed shaped tableaux is bounded from above, our proposition is proved. □

5. THE VARIETIES $\mathcal{F}^a(d_1, \dots, d_s)$

5.1. **The ideal I^a.** Throughout the section we fix a sequence $1 \le d_1 < \dots < d_s \le n-1$ and omit the numbers d_i in the notations $\mathcal{F}^a$, I^a, etc.

Let $U \hookrightarrow \mathcal{F}^a$ be the dense $\mathbb{G}_a^M$-orbit

$$U = \left\{ \exp\left(\sum_{1 \le i \le j \le n-1} c_{i,j} f_{i,j} \right) [v_{\omega_{d_1}}] \times \dots \times \exp\left(\sum_{1 \le i \le j \le n-1} c_{i,j} f_{i,j} \right) [v_{\omega_{d_s}}],\ c_{i,j} \in \mathbb{C} \right\}.$$

By definition, the variety $\mathcal{F}^a$ is the closure of U in the product of the projective spaces $\mathbb{P}(V^a_{\omega_{d_i}})$, $i = 1, \dots, s$. Recall the coordinates $X^a_{i_1,\dots,i_d}$ in $V^a_{\omega_d}$. These coordinates produce a multi-homogeneous coordinates on the product $\times_{i=1}^s \mathbb{P}(V^a_{\omega_{d_i}})$. Let us denote by $X^a_{i_1,\dots,i_d}(c_{i,j}) = X^a_{i_1,\dots,i_d}(\mathbf{c})$ the coordinates of a point

$$\exp\left(\sum_{1 \le i \le j \le n-1} c_{i,j} f_{i,j} \right) (v^a_{\omega_{d_1}} \times \dots \times v^a_{\omega_{d_s}}) \in \times_{i=1}^s V^a_{\omega_{d_i}}.$$

For instance, $X_{1,\dots,d} = 1$. Let us compute these coordinates explicitly. Let $1 \le r \le d$ be a number such that $i_r \le d$ and $i_{r+1} > d$. Define the numbers $1 \le j_1 < \dots < j_{d-r} \le d$ by the formula

$$\{j_1, \dots, j_{d-r}\} = \{1, \dots, d\} \setminus \{i_1, \dots, i_r\}.$$

Lemma 5.1. *We have*

$$X^a_{i_1,\dots,i_d}(\mathbf{c}) = (-1)^{\sum_{l=1}^r (i_l - l)} \sum_{\sigma\in S_{d-r}} (-1)^\sigma c_{j_1, i_{r+\sigma(1)}-1} \cdots c_{j_{d-r}, i_{r+\sigma(d-r)}-1}.$$

Proof. Lemma follows from formula (2.3). ☐

Corollary 5.2. *We have $X^a_{i_1,\dots,i_d}(\mathbf{c}) = \det C$, where $C_{\alpha,\beta}$ is an $n\times n$ matrix defined by*

$$C_{\alpha,\beta} = \begin{cases} c_{\alpha,i_\beta - 1}, & \text{if } \alpha \neq i_1,\dots,i_m, \beta > m, \\ 0, & \text{if } \alpha \neq i_1,\dots,i_m, \beta \le m, \\ 1, & \text{if } \alpha = i_1,\dots,i_m, \beta = \alpha, \\ 0, & \text{if } \alpha = i_1,\dots,i_m, \beta \neq \alpha. \end{cases} \tag{5.1}$$

We note that matrices (5.1) and (3.4) are identical up to the identification of parameters.

Example 5.3. Note that $X_{1,\dots,d}(\mathbf{c}) = 1$. Let $i \le d < j$. Then

$$X_{1,\dots,\hat{i}\dots,d,j}(\mathbf{c}) = (-1)^{d-i} c_{i,j-1}. \tag{5.2}$$

Lemma 5.4. *Fix a tuple $\mathbf{c}$. Then the coordinates $X^a_{i_1,\dots,i_{d_r}}(\mathbf{c})$, $r = 1,\dots,s$ satisfy the relations from the ideal I^a.*

Proof. Let $X_{i_1,\dots,i_d}(\mathbf{c})$ be the classical analogues of $X^a_{i_1,\dots,i_d}(\mathbf{c})$, i.e.

$$\exp\left(\sum_{1\le i\le j<n} c_{i,j} f_{i,j}\right) v_{\omega_d} = \sum_{1\le i_1<\dots<i_d\le n} X_{i_1,\dots,i_d}(\mathbf{c}) v_{i_1,\dots,i_d} \in V_{\omega_d}.$$

For any tuple $\mathbf{c}$ the coordinates $X_{i_1,\dots,i_d}(\mathbf{c})$, $d = d_1,\dots,d_s$ satisfy relations from I. We note that the polynomials $X^a_{i_1,\dots,i_{d_r}}(\mathbf{c})$ can be obtained from $X_{i_1,\dots,i_{d_r}}(\mathbf{c})$ simply by taking the lowest degree (in all variables $c_{i,j}$) terms. This lowest degree coincides with the PBW-degree of the tuple $i_1,\dots,i_d$. Since the terms in $R^{k;a}_{L,J}$ are exactly the lowest degree terms among those in $R^{k;a}_{L,J}$, the coordinates $X^a_{i_1,\dots,i_{d_r}}(\mathbf{c})$ do satisfy the relations $R^{k;a}_{L,J}$. ☐

Now consider the set of all zeros $X(I^a) \hookrightarrow \times_{r=1}^s \mathbb{P}(V^a_{\omega_{d_r}})$ of the ideal I^a.

Lemma 5.5. *Let $W \subset \times_{r=1}^s \mathbb{P}(V_{\omega_{d_r}})$ be an affine space defined by $X_{1,\dots,d_r} \neq 0$ for all $r = 1,\dots,s$. Then $X(I^a) \cap W = U$.*

Proof. Let $P_{i_1,\dots,i_{d_r}}$, $r = 1,\dots,s$ be the coordinates of a point $P \in X(I^a) \cap W$. We note that for all $l \le d_r < d_m < j$ the relation $R^{1;a}_{(1,\dots,d_m),(j,1,\dots,\hat{l}\dots,d_r)}$ give

$$P^a_{1,\dots,d_m} P^a_{j,1,\dots,\hat{l}\dots,d_r} = P^a_{j,1,\dots,\hat{l}\dots,d_m} P^a_{1,\dots,d_r}. \tag{5.3}$$

Assuming $P^a_{1,\dots,d} = 1$, relation (5.3) reads as

$$P^a_{1,\dots,\hat{l}\dots,d_m,j} = (-1)^{d_r+d_m} P^a_{1,\dots,\hat{l}\dots,d_r,j} \tag{5.4}$$

(which agrees with formula (5.2)). We set

$$c_{i,j-1} = (-1)^{d_m - i} P_{1,\dots,\hat{i}\dots,d_m,j},$$

if there exists m such that $i \le d_m < j$ and set $c_{i,j} = 0$ otherwise. Equation (5.4) guarantees that such tuple $c_{i,j}$ exists. Our goal is to show that

$$\exp\left(\sum_{1 \le i \le j < n} c_{i,j} f_{i,j}\right)\left([v^a_{\omega_{d_1}}] \times \dots \times [v^a_{\omega_{d_s}}]\right) = P. \tag{5.5}$$

The right and left hand sides belong to the affine space with the coordinates $X^a_{i_1,\dots,i_r}$ (we assume $X^a_{1,\dots,d_r} = 1$). We have defined $c_{i,j}$ in such a way that the coordinates $X^a_{1,\dots\hat{i},\dots,d_r,j}$ of the left and right hand sides of (5.5) coincide. This implies (5.5), since there is at most one point in $X(I^a) \cap W$ with the prescribed values of $X^a_{1,\dots\hat{i},\dots,d_r,j}$. In fact, let $J = (j_1, \dots, j_d)$ be a sequence such that $\deg J > 1$. Let us order J in such a way that $j_1 > d$. Then the relation $R^{1;a}_{(1,\dots,d),J}$ allows to rewrite X^a_J ($X^a_J X_{1,\dots,d}$ to be precise) as a polynomial in X^a_L with $\deg L < \deg J$. Now it suffices to note that the sequences $(1, \dots \hat{i}, \dots, d, j)$ are exactly the degree one sequences. $\square$

For a subset $R \subset \times_{r=1}^s \mathbb{P}(V^a_{\omega_{d_r}})$ let $I(R)$ be the ideal of multi-homogeneous polynomials vanishing on R. We now prove the main theorem of this section.

Theorem 5.6. *$I(U) = I^a$. Equivalently, $I(\mathcal{F}^a) = I^a$.*

Proof. We need to prove that if a multi-homogeneous polynomial F in variables $X^a_{i_1\dots i_d}$, $d = d_1, \dots, d_s$ vanishes on U, then $F \in I^a$. For a d-tuple $j_1, \dots, j_d$ consider the relation $R^{1;a}_{(1,\dots,d);(j_1,\dots,j_d)} \in I^a$. This relation is equal to

$$X^a_{1,\dots,d} X^a_{j_1,\dots,j_d} - \sum X^a_L X^a_{J'},$$

where for all terms $X^a_L X^a_{J'}$ we have

$$\deg X^a_L = 1, \ \deg X^a_{J'} = \deg(j_1, \dots, j_d) - 1.$$

(Recall the PBW degree $\deg J = \#\{l : j_l > d\}$). Therefore, the product $(X^a_{1,\dots,d})^{\deg J} X^a_J$ can be expressed modulo the ideal I^a as a homogeneous polynomial in coordinates $X^a_{1,\dots\hat{l}\dots,d,j}$ ($l \le d < j$).

We show that $F \in I^a$ in the following way: first assume that F depends only on the variables $X^a_{i_1,\dots,i_d}$ with a fixed d. Then F vanishes on a dense open cell of the Grassmanian $Gr(d, n)$ and hence the polynomial F belongs to I. Since $R^{k;a}_{L,J} = R^k_{L,J}$ (up to a superscript a) if lengths of L and J coincide, we obtain $F \in I^a$. Now consider the general case, i.e. assume that F depends on at least two groups of coordinates (with different d). Let r be the maximal number such that for all $m < r$ F does not depend on the variables $X^a_{i_1,\dots,i_{d_m}}$. We then show that there exists a polynomial G such that $G - F \in I^a$ and G is independent of the variables $X^a_{i_1,\dots,i_{d_m}}$ with $m \le r$.

Without loss of generality we assume that $r = 1$. First, there exists a number N such that the polynomial $\bar{F} = (X^a_{1,\dots,d_1})^N F$ can be expressed modulo I^a as a polynomial independent of $X^a_{j_1,\dots,j_d}$ whenever $\deg(j_1, \dots, j_d) > 1$. Note also that $\bar{F}$ still vanishes on U. Recall the expression (5.2) for the degree one coordinates:

$$X^a_{1,\dots\hat{l}\dots,d_1,m}(\mathbf{c}) = (-1)^{d_1-l} c_{l,m-1}.$$

From this formula and since $\bar{F}$ vanishes on the whole U, $\bar{F}$ does not depend on $X^a_{1,\dots\hat{l}\dots,d_1,m}$ with $m \le d_2$ (because numbers $c_{l,m-1}$ with $m \le d_2$ can come nowhere, but from $X^a_{1,\dots\hat{l}\dots,d_1,m}(\mathbf{c})$). For the rest $X^a_{1,\dots\hat{l}\dots,d_1,m}$ we have the following relation in I^a:

$$X^a_{1,\dots,d_2} X^a_{1,\dots\hat{l}\dots,d_1,m} - (-1)^{d_1+d_2} X^a_{1,\dots\hat{l},\dots,d_2,m} X^a_{1,\dots,d_1} = 0.$$

Therefore, there exist numbers M_1 and M_2 and a polynomial G such that

$$(X^a_{1,\dots,d_2})^{M_1} \bar{F} - (X^a_{1,\dots,d_1})^{M_2} G \in I^a$$

and G does not depend on variables $X^a_{i_1,\dots,i_{d_1}}$. Note that since $\bar{F}$ vanishes on U and $X^a_{1,\dots,d}$ equals to 1 on U, the polynomial G vanishes on U as well. By induction on s we can assume $G \in I^a$. Therefore,

$$(X^a_{1,\dots,d_2})^{M_1} \bar{F} = (X^a_{1,\dots,d_2})^{M_1} (X^a_{1,\dots,d_1})^N F \in I^a.$$

Since I^a is prime (see Corollary 4.11), we arrive at $F \in I^a$. □

5.2. **Flatness.** We close with the proof of the flatness of the degeneration $\mathcal{F} \to \mathcal{F}^a$. Let t be a variable. We define an algebra Q^t over the ring $\mathbb{C}[t]$ as a quotient of the polynomial ring $\mathbb{C}[t][X_{i_1,\dots,i_d}]$, $d = d_1, \dots, d_s$ by the ideal I^t generated by quadratic relations $R^{k;t}_{L,J}$. These relations are t-deformations of the relations $R^k_{L,J}$. Namely, let $R^k_{L,J} = \sum_i X_{L^{(i)}} X_{J^{(i)}}$. Then

$$R^{k;t}_{L,J} = t^{-\min_i(\deg L^{(i)} + \deg J^{(i)})} \sum_i t^{\deg L^{(i)} + \deg J^{(i)}} X_{L^{(i)}} X_{J^{(i)}}.$$

The following lemma describes the fibers of Q^t.

Lemma 5.7. *$Q^t/(t) \simeq Q^a$, $Q^t/(t-u) \simeq Q$ for $u \neq 0$.*

Proof. Straightforward. □

Proposition 5.8. *Q^t is $\mathbb{C}[t]$ free.*

Proof. We prove that the elements X_T, $T \in ST^a_\lambda$, $\lambda \in P^+(d_1, \dots, d_s)$ form a $\mathbb{C}[t]$ basis of Q^t. First, let us prove that the elements X_T are linearly independent. Assume

$$\sum_T m_T X_T = \sum_i R^{k_i;t}_{L^{(i)},J^{(i)}} p_i(t), \ m_T, p_i \in \mathbb{C}[t].$$

There exists a number $u \in \mathbb{C}$ such that $m_T(u) \neq 0$ if $m_T \neq 0$. Now Proposition 4.12 gives linear independence. The proof of the spanning property is analogous to the proof of Proposition 4.12. □

Acknowledgments

We are grateful to I.Arzhantsev, M.Finkelberg and A.Kuznetsov for useful discussions and to I.Arzhantsev for useful remarks on the earlier version of the paper. This work was partially supported by the Russian President Grant MK-281.2009.1, RFBR Grants 09-01-00058, 07-02-00799 and NSh-3472.2008.2, by Pierre Deligne fund based on his 2004 Balzan prize in mathematics and by EADS foundation chair in mathematics.

References

[A] I. Arzhantsev, *Flag varieties as equivariant compactifications of* $\mathbb{G}_a^n$, arXiv:1003.2358.

[AS] I. Arzhantsev, E. Sharoiko, *Hassett-Tschinkel correspondence: modality and projective hypersurfaces*, arXiv:0912.1474.

[C] P. Caldero, *Toric degenerations of Schubert varieties*, Transformation Groups, Vol. 7, No. 1, (2002) 51-60.

[Fe1] E. Feigin, *The PBW filtration*, Represent. Theory 13 (2009), 165-181.

[Fe2] E. Feigin, *The PBW filtration, Demazure modules and toroidal current algebras*, SIGMA, 4 (2008), 070, 21 p.

[Fu] W. Fulton, *Young tableaux, with applications to representation theory and geometry.* Cambridge University Press, 1997.

[FH] W. Fulton, J. Harris, (1991), *Representation theory. A first course*, Graduate Texts in Mathematics, Readings in Mathematics, 129, New York.

[FFL] B. Feigin, E. Feigin, P. Littelmann, *Zhu's algebras,* C_2*-algebras and abelian radicals*, arXiv:0907.3962.

[FFoL] E. Feigin, G. Fourier, P. Littelmann, *PBW filtration and bases for irreducible modules in type* A_n, arXiv:1002.0674.

[FL] E. Feigin, P. Littelmann, *Zhu's algebra and the* C_2*-algebra in the symplectic and the orthogonal cases*,2010 J. Phys. A: Math. Theor. 43 135206.

[GL] N. Gonciulea, V. Lakshmibai, *Degenerations of flag and Schubert varieties to toric varieties*, Transformation Groups, vol 1, no:3 (1996), 215-248.

[HT] B. Hassett, Yu. Tschinkel, *Geometry of equivariant compactifications of* $\mathbb{G}_a^n$, Int. Math. Res. Notices 20 (1999), 1211-1230.

[K1] S. Kumar, *Kac-Moody groups, their flag varieties and representation theory,* Progress in Mathematics vol. 204, Birkhauser, Boston, 606 (2002).

[K2] S. Kumar, *The nil Hecke ring and singularity of Schubert varieties*, Inventiones Math. 123, 471-506 (1996).

[L] V. Lakshmibai, *Degenerations of flag varieties to toric varieties*, C. R. Acad. Sci. Paris, 321 (1995), 1229-1234.

[V] E. Vinberg, *On some canonical bases of representation spaces of simple Lie algebras*, conference talk, Bielefeld, 2005.

Evgeny Feigin:
Tamm Theory Division, Lebedev Physics Institute,
Leninisky prospect, 53, 119991, Moscow, Russia,
and
French-Russian Poncelet Laboratory, Independent University of Moscow
E-mail address: `evgfeig@gmail.com`